# **QUADRATIC FUNCTIONS**

### **A DEFINITION**

Definition Quadratic Function -


A quadratic function is  $x \mapsto ax^2 + bx + c$  where  $a \neq 0$ .

Ex: For  $f(x) = x^2 - 3x + 1$ , evaluate f(2).

Answer: 
$$f(2) = (2)^2 - 3(2) + 1$$
  
=  $4 - 6 + 1$   
=  $-1$ 


#### **B GRAPH**

The parabola is one of the conic sections, which are the group of curves obtained by intersecting a cone with a plane. A parabola is produced by intersecting the cone with a plane parallel to its generating line. By intersecting the cone at other angles, we can produce circles, hyperbolas, and ellipses.



#### Definition Parabola


Given a quadratic function  $x \mapsto ax^2 + bx + c$  where  $a \neq 0$ , its graph is called a parabola.



**Ex:** Sketch the graph of  $x \mapsto x^2 - x - 1$ .

Answer: A table of values is

$\boldsymbol{x}$	-2	-1	0	1	2
y	5	1	-1	-1	1



To understand the concavity based on the sign of a, use the GeoGebra animation at https://www.geogebra.org/m/gn3c2sqe.

### Proposition Concavity -

For any quadratic function  $x \mapsto ax^2 + bx + c$ ,  $a \neq 0$ :

- If a > 0, the graph is concave up:
- If a < 0, the graph is concave down:

## C SOLVING f(x) = y

#### Method **Solving** f(x) = y

When solving for a value of f(x) = y, we obtain a quadratic equation in x. Since it is quadratic, there may be 0, 1, or 2 real solutions for x.

Ex: For  $f(x) = 2x^2 - 5x + 2$ , find the x-intercepts of the function.

Answer: Set f(x) = 0:  $2x^2 - 5x + 2 = 0$ , with a = 2, b = -5, c = 2.

• 
$$\Delta = b^2 - 4ac$$
  
=  $(-5)^2 - 4(2)(2)$   
=  $25 - 16$   
=  $9$ 

• As  $\Delta > 0$ , there are 2 distinct roots.

$$\bullet \ x = \frac{-b - \sqrt{\Delta}}{2a} \qquad \text{or } x = \frac{-b + \sqrt{\Delta}}{2a}$$

$$x = \frac{-(-5) - \sqrt{9}}{2 \cdot 2} \qquad \text{or } x = \frac{-(-5) + \sqrt{9}}{2 \cdot 2}$$

$$x = \frac{5 - 3}{4} \qquad \text{or } x = \frac{5 + 3}{4}$$

$$x = \frac{2}{4} \qquad \text{or } x = \frac{8}{4}$$


$$x = \frac{1}{2} \qquad \text{or } x = 2$$

The x-intercepts are at  $x = \frac{1}{2}$  and x = 2.


## Proposition Relative position between the graph and the x-axis

For any quadratic function  $x\mapsto ax^2+bx+c$  and the discriminant  $\Delta=b^2-4ac$ :

• If  $\Delta > 0$ , then the graph intersects the x-axis twice.



• If  $\Delta = 0$ , then the graph touches the x-axis at one point.



• If  $\Delta < 0$ , then the graph does not intersect the x-axis.

